

## IDH2(R172H)

**Catalog Number:** 26084

**Gene Symbol:** IDH2(R172H)

**Description:** Anti-IDH2(R172H) Mouse Monoclonal Antibody

**Background:** Isocitrate dehydrogenase (IDH) catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. The isocitrate and isopropylmalate dehydrogenases family has three members, IDH1, IDH2 and IDH3. IDH2 plays a role in intermediary metabolism and energy production. Defects in IDH2 are the cause of D-2-hydroxyglutaric aciduria type 2 (D2HGA2). Somatic mosaic mutations of this protein have also been found associated to Ollier disease and Maffucci syndrome., and R172H IDH2 mutations do exist in diffusely infiltrative gliomas.

**Immunogen:** A synthetic peptide from the internal region of IDH2 which includes the mutation of R172H, human origin.

**Tested applications:** ELISA, WB, IHC

**Recommended dilutions:**

ELISA: 1:1000-1:5000

WB: 1:100-1:1000

IHC: 1:50-1:100

IF: 1:50-1:100

**Concentration:** 1 mg/ml

**Host:** Mouse

**Clonality:** Monoclonal

**Purity:** Purified from ascites

**Format:** Liquid

**Storage buffer:**


Preservative: no

Constituents: PBS (without  $Mg^{2+}$  and  $Ca^{2+}$ ), pH 7.4, 150 mM NaCl, 50% glycerol


**Species Reactivity:** Anti-IDH2(R172H) antibody recognizes IDH2(R172H) of vertebrates.

**Storage Conditions:** Store at -20 °C. Avoid freeze / thaw cycles.

**Western blot:**



**Immunofluorescence:**



**Immunofluorescence of cells expressing IDH2 proteins with anti-IDH2 (R172H) antibody.**

HEK293T cells were transfected with pCDNA3-GFP-IDH2 (WT) plasmid (left column) or pCDNA3-GFP-IDH2 (R172H) plasmid (right column), then fixed and stained with anti-IDH2 (R172H) monoclonal antibody (Cat. #26084).